arXiv:1304.1695v3 [math.AG] 7 May 2013

DEFORMING GEOMETRIC TRANSITIONS

MICHELE ROSSI

Dedicated to Professor D. Gallarati on the occasion of his 90th birthday

Abstract. After a quick review of the wild structure of the complex moduli space of Calabi–Yau threefolds and the role of geometric transitions in this context (the Calabi–Yau web) the concept of *deformation equivalence* for geometric transitions is introduced to understand the arrows of the Gross-Reid Calabi–Yau web as deformation-equivalence classes of geometric transitions. Then the focus will be on some results and suitable examples to understand under which conditions it is possible to get *simple* geometric transitions.

Contents

Introduction 2
1. Calabi–Yau varieties 2
2. Aspects of deformation theory of Calabi–Yau varieties 4
2.1. Deformation equivalence of Calabi–Yau varieties 5
3. Geometric transitions 5
3.1. The basic example: the conifold transition in $\mathbb{P}^4$ 6
3.2. Local topology of a conifold transition 7
4. The Reid's fantasy 8
5. The string theoretic Calabi–Yau web 8
5.1. The *vacuum degeneracy problem* 9
6. The Gross Calabi–Yau web: nodes and arrows 9
7. Deformation of a morphism 11
7.1. Deformation equivalence of geometric transitions 11
8. Simple geometric transitions 13
8.1. Type II geometric transitions are not simple 13
8.2. An example of a non-simple small geometric transition 14
8.3. A cohomological condition for simplicity of small transitions 15
8.4. An example of a simple small geometric transition 16
References 18

This research was partially supported by the MIUR-PRIN 2010-11 "Geometria delle Varietà Algebriche" Research Funds.

INTRODUCTION

These notes were written on the occasion of the GTM Seminar held in Genova on March 21-22, 2013, and dedicated to Prof. D. Gallarati for his 90th birthday.

First of all I would like to thank the organizers A. Alzati, G. Casnati, F. Galluzzi, R. Notari, M.E. Rossi, G. Valla and, last but not least, L. Badescu and M. Beltrametti, for the warm welcome and hospitality and all the participants for the pleasant climate they were able to establish during the two days of work. Among them a special thanks to F. Catanese for stimulating conversations and suggestions. Finally I would like to thank R. Davies for his helpful comments.

The aim of this paper is that of quickly speaking about the moduli space of Calabi–Yau varieties, or better, the Calabi–Yau web in its different formulations, both in mathematics and in physics.
A large first part of these notes (sections from 1 to 6) has a purely expository purpose. For a broader discussion of these aspects the interested reader is referred to the extensive survey on the subject [31]. A second part (sections from 7 to 8) is instead devoted to giving some new ideas, partial results and examples, with the aim of shedding a brighter light on the study of geometric transitions and, more generally, on the Calabi–Yau moduli space. In particular, the notion of *deformation equivalence of geometric transitions* is introduced (see Definition 7.1), which allows us, on the one hand, to think of the Gross-Reid Calabi–Yau web as a kind of quotient of the string theoretic Calabi–Yau web by means of def-equivalence, and, on the other hand, to isolate a class of geometric transitions (referred to as *simple*, see Definition 8.1) having the properties of being well understood both from the physical and the mathematical point of view, guaranteeing the existence of an *almost diffeomorphism* (see Remark 8.2) between the Calabi–Yau threefolds they connect. New results in this context are then given by Proposition 7.3 and Theorem 8.5. The former gives an easier formulation of def-equivalence between geometric transitions admitting singular loci comprising at most isolated terminal singularities; this is the case e.g. for *small* geometric transitions (see Definition 8.6). The latter characterizes the *type II* geometric transitions (see Definition 8.3) as *never simple* ones. This paper ends up by studying when a small geometric transition is actually a simple one: examples of both simple and non-simple small geometric transitions are given. The main result in this context is given by Theorem 8.8, which gives a *generally sufficient* cohomological condition for small geometric transitions to be simple: here the generality hypothesis is quite strong, hiding technical details whose explanation is beyond the scope of this article. The interested reader is referred to the forthcoming paper [33] for a full proof.

1. CALABI–YAU VARIETIES

Definition 1.1. A compact, complex, Kähler manifold Y is a *Calabi–Yau variety* if

(1) $\bigwedge^n \Omega_Y =: \mathcal{K}_Y \cong \mathcal{O}_Y$
(2) $h^{p,0}(Y) = 0 \quad \forall 0 < p < \dim Y$

A n–dimensional Calabi–Yau variety will be also called a *Calabi–Yau n–fold*.

Remarks 1.2. (1) The given definition of Calabi–Yau variety includes the following lower dimensional cases
- smooth elliptic curves,

  - smooth $K3$ surfaces.

(2) Observe that although a smooth elliptic curve always admits a projective embedding, this is no longer the case for $K3$ surfaces. By the way, for $\dim Y \geq 3$, the given definition of a Calabi–Yau variety Y implies that Y is a *projective variety*: the embedding can be fixed by a suitable integer multiple of a rational Kähler form near enough to the Kähler metric of Y.

Examples 1.3. (1) Smooth hypersurfaces of degree $n+1$ in $\mathbb{P}^n$ (use Adjunction Formula and the Lefschetz Hyperplane Theorem).

(2) Smooth hypersurfaces (if exist!) of a weighted projective space $\mathbb{P}(q_0, \ldots, q_n)$ of degree $d = \sum_{i=0}^{n} q_i$.

(3) The general element of the anti–canonical system of a *sufficiently good* 4–dimensional toric Fano variety (see [1]).

(4) Suitable complete intersections.... (iterate the previous examples).

(5) The double covering of $\mathbb{P}^3$ ramified along a smooth surface of degree 8 in $\mathbb{P}^3$ (octic double solid).

In dimension greater than or equal to 3 the previous examples give *topologically distinct* complex varieties, implying immediately that *the complex moduli space of Calabi–Yau n–folds, with $n \geq 3$, has to be necessarily disconnected.* This fact apparently clashes with the smaller dimensional cases:

- the complex moduli space of elliptic curves is given by the modular curve $\Gamma(1)\backslash\mathbb{H} \cong \mathbb{P}^1$ which parameterizes complex structures over the topological torus $S^1 \times S^1$,
- after Kodaira [20], the complex moduli space of $K3$ surfaces is given by a *smooth, complex, irreducible space of dimension 20.*

Anyway if we insist on looking at this situation from the algebraic point of view, then the moduli space of algebraic $K3$ surfaces turns out to be a dramatically more complicated object: the following facts were known to F. Enriques [11]:

- $\forall g \geq 3$ there exists a K3 surface of degree $2g-2$ in $\mathbb{P}^g$; hence its sectional genus is g;
- $\forall g \geq 3$ we can obtain a space $\mathcal{M}_g$ of complex projective moduli of such surfaces, by imposing a polarization: $\mathcal{M}_g$ is an irreducible, analytic variety with $\dim_{\mathbb{C}} \mathcal{M}_g = 19$;
- then the complex moduli space $\mathcal{M}^{alg}$ of algebraic K3 surfaces is a *reducible* analytic variety and it admits a countable number of irreducible components;
- there exist K3 surfaces belonging to more than one irreducible component of $\mathcal{M}^{alg}$; anyway if we restrict to K3's admitting Pic $\cong \mathbb{Z}$ (they give the general element of any irreducible component) then they belong to only one irreducible component.

What could appear to F. Enriques as a wildly reducible moduli space was explained by K. Kodaira [20] as an analytic codimension 1 subvariety of a smooth, irreducible, analytic variety $\mathcal{M}$. More precisely:

- there exist analytic non-algebraic K3 surfaces,
- the Kuranishi space of any analytic K3 surface is smooth and of dimension 20.

The latter suffices to construct a smooth, irreducible, analytic universal family of K3 surfaces: its base $\mathcal{M}$ is the complex analytic moduli space of K3 surfaces and $\dim_{\mathbb{C}} \mathcal{M} = 20$. Moreover $\mathcal{M}^{alg}$ turns out to be a dense subset of $\mathcal{M}$.

In other terms, Kodaira recovered an irreducible moduli space for $K3$'s by leaving the algebraic category and working in the bigger category of compact complex surfaces. By this observation, in the late 80's, M.Reid [30] proposed a conjectural construction of a *sort of connected moduli space for Calabi–Yau 3-folds* suggesting a construction (originally due to F. Hirzebruch and later called *conifold transition*) to parameterize birational classes of Calabi–Yau 3-folds by means of moduli of complex structures on suitable non-Kähler complex 3-folds given by the connected sum of copies of solid hypertori $S^3 \times S^3$: this is the famous *Reid's fantasy*.

2. Aspects of deformation theory of Calabi–Yau varieties

Let $\mathcal{X} \xrightarrow{f} B$ be a *flat*, surjective map of complex spaces such that B is connected and there exists a special point $o \in B$ whose fibre $X = f^{-1}(o)$ may be singular. Then $\mathcal{X}$ is called *a deformation family of* X. If the fibre $X_b = f^{-1}(b)$ is smooth, for some $b \in B$, then X_b is called *a smoothing of* X. Moreover X_b is also called a *deformation* of X_o.

If the morphism f is *smooth* then $\mathcal{X} \xrightarrow{f} B$ is called a *smooth deformation family.*

Given a deformation family $\mathcal{X} \xrightarrow{f} B$ of X *for each point* $b \in B$ *there is a well defined linear (and functorial) map*

$$D_b f : \ T_b B \longrightarrow \mathbb{T}^1_{X_b} := \operatorname{Ext}^1\left(\Omega^1_X, \mathcal{O}_X\right) \qquad \text{(Generalized Kodaira–Spencer map)}$$

(see e.g. [26] Theorem 5.1). Recall that $\mathcal{X} \xrightarrow{f} B$ is called

- a *versal* deformation family of X if for any deformation family $(\mathcal{Y}, X) \xrightarrow{g} (C, 0)$ of X there exists a map of pointed complex spaces $h : (U, 0) \to (B, o)$, defined on a neighborhood $0 \in U \subset C$, such that $\mathcal{Y}|_U$ is the *pull–back* of $\mathcal{X}$ by h i.e.
$$\begin{tikzcd}
 & \mathcal{Y}|_U = U \times_B \mathcal{X} \arrow[r] \arrow[d, "g"] & \mathcal{X} \arrow[d, "f"] \\
C & U \arrow[l, hook] \arrow[r, "h"] & B
\end{tikzcd}$$
- an *effective versal* deformation family of X if it is versal and the generalized Kodaira–Spencer map evaluated at $o \in B$, $D_o f : \ T_o B \longrightarrow \mathbb{T}^1_X$ is injective,
- a *universal* family if it is versal and h is defined on every point of C and is uniquely determined in a neighborhood $0 \in U \subset C$.

Theorem 2.1 (Douady–Grauert–Palamodov [10], [16], [25] and [26] Theorems 5.4-6)**.** *Every compact complex space* X *has an effective versal deformation* $\mathcal{X} \xrightarrow{f} B$ *which is a proper map and a versal deformation of each of its fibers. Moreover the germ of analytic space* (B, o) *is isomorphic to the germ of analytic space* $(q^{-1}(0), 0)$*, where* $\mathbb{T}^i(X) := \operatorname{Ext}^i\left(\Omega^1_X, \mathcal{O}_X\right)$ *and* $q : \mathbb{T}^1_X \to \mathbb{T}^2_X$ *is a suitable holomorphic map (the* obstruction map*) such that* $q(0) = 0$*. In particular if* $\mathbb{T}^0_X = 0$ *then the previous versal effective deformation of* X *is actually a universal one for all the fibres close enough to* X*.*

Since for a Calabi–Yau variety Y we get $\mathbb{T}^0(Y) \cong H^0(Y, \Theta_Y) \cong H^0(Y, \Omega_Y^{n-1}) = 0$ then

Corollary 2.2. *Every Calabi–Yau variety* Y *admits a universal effective family of Calabi–Yau deformations of* Y*. In particular* $h^{n-1,1}(Y)$ *turns out to be the dimension of the complex moduli space of* Y

Definition 2.3 (Kuranishi space)**.** The germ of analytic space

$$\mathrm{Def}(X) := (B, o)$$

defined in the previous Theorem, is called the *Kuranishi space of* X.

Theorem 2.4 (Bogomolov-Tian-Todorov [3],[36],[37],[28])**.** *Any Calabi–Yau variety have unobstructed deformations, i.e., its Kuranishi space is smooth.*

2.1. **Deformation equivalence of Calabi–Yau varieties.** In the late 80's R. Friedman and J.W. Morgan [14] introduced the following equivalence relation between complex manifolds. Here we use notation introduced by F. Catanese and M. Manetti in several subsequent discussions of related problems and conjectures [4],[5],[22].

Definition 2.5 (Deformation equivalence, [14] pg. 10)**.** Two smooth complex manifolds X_1 and X_2 are *deformation equivalent* (i.e. either *def-equivalent* or *have the same deformation type*) if there exists a smooth deformation family $\mathcal{X} \xrightarrow{f} B$ whose base B is a connected complex space admitting two points $b_1, b_2 \in B$ such that

$$X_i = f^{-1}(b_i) \ , \ i = 1, 2 \ .$$

We will write $X_1 \sim X_2$. This gives an equivalence relation on the set of complex manifolds.

By the classical theorem of Ehresmann:

- *two def-equivalent complex manifolds are orientedly diffeomorphic.*

Freidman and Morgan conjectured that the converse could be true: this is the so called *def = diff problem.* Friedman actually proved this equivalence for *compact complex surfaces with* $b_1 = 0$ *and Kodaira dimension less than or equal to 1* [13]: in particular *it holds for* $K3$ *surfaces.*

However the general problem has been negatively settled by several counterexamples, the first of which was given by M. Manetti, then followed by many others by F. Catanese, Kharlamov-Kulikov, Bauer-Catanese-Grunewald, Catanese-Wajnryb ... (see [2] and therein references).

Anyway there is a point which is worth to be emphasized: as far as I know, all these counterexamples are given by surfaces (and/or 4-manifolds), leaving completely open the the higher dimensional problem. In particular:

- *since the def = diff problem admits a positive answer for elliptic curves (obvious) and* $K3$ *surfaces (Friedman* [13]*) it makes sense to ask if does it hold for any Calabi–Yau varieties.*

3. Geometric transitions

Definition 3.1. Let Y be a Calabi–Yau n–fold and $\phi : Y \to \overline{Y}$ be a *birational contraction* onto a *normal* variety. If there exists a complex deformation (*smoothing*) of $\overline{Y}$ to a Calabi–Yau n–fold $\widetilde{Y}$, then the process of going from Y to $\widetilde{Y}$ is called a *geometric transition* (for short *g.t.*) and denoted by $T(Y,\overline{Y},\widetilde{Y})$ or by the diagram

$$\begin{tikzcd} Y \arrow[r, "\phi"] \arrow[rr, bend right, dotted, "T"'] & \overline{Y} \arrow[r, leftrightsquigarrow, dashed] & \widetilde{Y} \end{tikzcd} .$$

A g.t. $T(Y,\overline{Y},\widetilde{Y})$ is called *trivial* if $\widetilde{Y}$ is a deformation of Y.
A g.t. $T(Y,\overline{Y},\widetilde{Y})$ is called a *conifold transition* (for short *c.t.*) if $\overline{Y}$ admits only *ordinary double points* (nodes) as singularities.

Remarks 3.2.
(1) *Trivial g.t.s may occur*: in fact it is not possible to realize non–trivial transitions in dimension less than or equal to 1. For a trivial g.t. in dimension 3 one may e.g. consider Example 4.6 in [39] where ϕ admits an elliptic scroll as exceptional divisor and contracts it down to an elliptic curve C.
(2) The transition process was firstly (locally) observed by H. Clemens in the study of double solids V admitting at worst nodal singularities [7]: in his Lemma 1.11 he pointed out “the relation of the resolution of the singularities of V to the standard $S^3 \times D_3$ to $S^2 \times D_4$ surgery”.

3.1. **The basic example: the conifold transition in $\mathbb{P}^4$.** The following example, given in [17], shows that *non–trivial (conifold) transitions occur when* $\dim Y \geq 3$.

Let $\overline{Y} \subset \mathbb{P}^4$ be the singular hypersurface given by the following equation

$$x_3 g(x_0,\dots,x_4) + x_4 h(x_0,\dots,x_4) = 0 \tag{1}$$

where g and h are generic homogeneous polynomials of degree 4. $\overline{Y}$ is then the *generic quintic 3–fold containing the plane* $\pi : x_3 = x_4 = 0$. Then the singular locus of $\overline{Y}$ is given by

$$\mathrm{Sing}(\overline{Y}) = \{[x] \in \mathbb{P}^4 | x_3 = x_4 = g(x) = h(x) = 0\} . \tag{2}$$

One can then easily prove that:

(1) $\mathrm{Sing}(\overline{Y})$ is composed by 16 nodes,
(2) *(the resolution Y)*: $\mathrm{Sing}(\overline{Y})$ can be simultaneously resolved and the resolution $\phi : Y \to \overline{Y}$ is a *small blow up* such that Y is a smooth Calabi–Yau 3–fold,
(3) *(the smoothing $\widetilde{Y}$)*: $\overline{Y}$ admits the obvious smoothing given by the generic quintic 3–fold $\widetilde{Y} \subset \mathbb{P}^4$. In particular $\widetilde{Y}$ cannot be a deformation of Y i.e. the conifold transition $T(Y,\overline{Y},\widetilde{Y})$ is not trivial.

The latter fact can be easily shown by applying the Lefschetz Hyperplane Theorem and the Künneth Formula to get the following relations on the second Betti numbers:

$$\begin{aligned} b_2(\widetilde{Y}) &= b_2(\mathbb{P}^4) = 1 \\ b_2(Y) &= b_2(\mathbb{P}^4 \times \mathbb{P}^1) = 2 \end{aligned} \tag{3}$$

Therefore $\widetilde{Y}$ and Y cannot be smooth fibers of the same analytic family.

3.2. **Local topology of a conifold transition.** From now on we will restrict to consider the case $n = 3$ of Calabi–Yau 3-folds. Then we can observe the following facts (for full details the interested reader is referred to [15], §1.1).

(1) Locally a 3-dimensional node can be described by the local equation
$$\overline{U} := \{z_1 z_3 + z_2 z_4 = 0\} \subset \mathbb{C}^4 \ .$$
Topologically $\overline{U}$ turns out to be a cone over $S^3 \times S^2$.

(2) A local resolution of $\overline{U}$ is described by
$$\widehat{U} := \left\{ \begin{array}{lcl} y_0 z_4 - y_1 z_3 & = & 0 \\ y_0 z_1 + y_1 z_2 & = & 0 \end{array} \right\} \subset \mathbb{C}^4 \times \mathbb{P}^1 \ .$$
Then there exists a diffeomorphism $\widehat{U} \overset{\Phi}{\cong} \mathbb{R}^4 \times S^2$. Moreover $\widehat{U}$ can be identified with the total space of the rank 2 holomorphic vector bundle $\mathcal{O}_{\mathbb{P}^1}(-1) \oplus \mathcal{O}_{\mathbb{P}^1}(-1)$ over the exceptional fibre $\mathbb{P}^1_{\mathbb{C}} = \varphi^{-1}(0)$. In particular $\widehat{U}$ admits a natural complex structure.

(3) A local smoothing $\widetilde{U}$ of the node $\overline{U}$ can be given by the 1–parameter family $f : \mathcal{U} \to \mathbb{R}$ where
$$U_t := f^{-1}(t) = \{z_1 z_3 + z_2 z_4 = t\} \subset \mathbb{C}^4 \ .$$
Setting $\widetilde{U} := U_{t_0}$ for some $t_0 \in \mathbb{R}, t_0 > 0$, then $\widetilde{U} \overset{\Psi}{\cong} S^3 \times \mathbb{R}^3$ since it is diffeomorphic to the cotangent bundle T^*S^3 of the 3–sphere giving the vanishing cycle of the smoothing. In particular $\widetilde{U}$ admits a natural symplectic structure for which the vanishing sphere turns out to be a lagrangian submanifold.

Theorem 3.3 (Clemens [7] Lemma 1.11, [15] Thm. 1.6)**.** *Let $D_n \subset \mathbb{R}^n$ be the closed unit ball and consider*

- $S^3 \times D_3 \subset S^3 \times \mathbb{R}^3 \overset{\Psi^{-1}}{\cong} \widetilde{U}$
- $D_4 \times S^2 \subset \mathbb{R}^4 \times S^2 \overset{\Phi^{-1}}{\cong} \widehat{U}$

Then $\widetilde{D} := \Psi^{-1}(S^3 \times D_3)$ and $\widehat{D} := \Phi^{-1}(D_4 \times S^2)$ are compact tubular neighborhoods of the vanishing cycle $\widetilde{S} \subset \widetilde{U}$ and of the exceptional cycle $\mathbb{P}^1_{\mathbb{C}} \subset \widehat{U}$, respectively. Consider the standard diffeomorphism
$$\begin{array}{cccc} \alpha' : & (\mathbb{R}^4 \setminus \{0\}) \times S^2 & \overset{\cong}{\longrightarrow} & S^3 \times (\mathbb{R}^3 \setminus \{0\}) \\ & (u, v) & \longmapsto & (\frac{u}{|u|}, |u| v) \end{array}$$
and restrict it to $D_4 \times S^2$. Since
$$\partial(D_4 \times S^2) = S^3 \times S^2 = \partial(S^3 \times D_3)$$
observe that $\alpha'|_{\partial(D_4 \times S^2)} = \mathrm{id}|_{S^3 \times S^2}$. Hence α' induces a standard surgery from $\mathbb{R}^4 \times S^2$ to $S^3 \times \mathbb{R}^3$.
Then $\widetilde{U}$ can be obtained from $\widehat{U}$ by removing $\widehat{D}$ and pasting in $\widetilde{D}$, by means of the diffeomorphism $\alpha := \Psi^{-1} \circ \alpha' \circ \Phi$.

Let us underline a global consequence of the Theorem 3.3, as a straightforward application of the Seifert-van Kampen Theorem

Corollary 3.4. *A conifold transition does not change the fundamental group.*

4. The Reid's fantasy

Since g.t.s (and in particular *conifold* t.s) may connect topologically distinct Calabi–Yau 3-folds, M. Reid thought that they could be the right instrument to recover a sort of *connectedness* of the Calabi–Yau 3-folds complex moduli space [30]. Quickly, his construction was the following.

1. *Assumption*: every projective Calabi–Yau 3–fold Y is birational to a Calabi–Yau 3–fold Y' such that $H^2(Y')$ is generated by rational curves.
2. Consequently if $\phi : Y' \to \overline{Y}$ is the morphism contacting all of them, then $\overline{Y}$ is always smoothable, by a Friedman result ([12], Corollary 4.7), and every smoothing $\widetilde{Y}$ has $b_2(\widetilde{Y}) = 0$, meaning that $\overline{Y}$ can be smoothed only to *non–Kähler* compact complex 3–folds.
3. By results of C. T. C. Wall [40], any such smoothing $\widetilde{Y}$ has topological type completely determined by its third Betti number $b_3(\widetilde{Y})$, implying that it is diffeomorphic to a connected sum $\left(S^3 \times S^3\right)^{\# r}$ of r copies of the *solid hypertorus* $S^3 \times S^3$.

Then we get the famous:

Conjecture 4.1 (the Reid's fantasy)**.** *Up to some kind of inductive limit over* r*, the birational classes of projective Calabi–Yau 3–folds can be fitted together, by means of geometric transitions, into one irreducible family parameterized by the moduli space* $\mathcal{N}$ *of complex structures over suitable connected sum of copies of solid hypertori.*

This conjecture has the further fascinating property of recovering the idea that moduli could be described by studying complex structures over a (hyper)-torus, typical of elliptic curves.
Unfortunately, the description of the moduli space $\mathcal{N}$ turns out to be a quite hard problem.

5. The string theoretic Calabi–Yau web

Calabi–Yau 3–folds play a fundamental role in 10–dimensional string theories: locally, four dimensions give rise to the usual Minkowsky space–time M_4 while the remaining six dimensions (the so called *hidden dimensions* for their microscopic extension) are compactified to a geometric model Y which, essentially to preserve the required supersymmetry, turns out to be a Calabi–Yau 3–fold. Therefore the string theoretic space-time looks like

- *a locally trivial 10-dimensional bundle whose base is the usual space-time of Einstein and which is locally isomorphic to* $M_4 \times Y$.

5.1. **The *vacuum degeneracy problem.*** In spite of the fact that there are only five consistent 10–dimensional super–string theories, actually nearly unique via dualities,



the compactification process give rise to the problem of choosing the appropriate Calabi–Yau model: which fibers in the space-time bundle?

In fact on the physical side, there is not any prescription for making a precise choice of the *vacuum model* and on the mathematical side there is a multitude of topologically distinct Calabi–Yau 3–folds. By the way, the choice of two distinct models does not at all give rise to equivalent physical theories, since the physics turns out to be strictly related to the cohomology of the Calabi–Yau model.

Ideas connected with the formulation of the Reid's fantasy, suggested to physicists, such as P. Candelas, P. S. Green, T. Hübsch and others that:

- *(simply connected) Calabi–Yau 3–folds could be, at least mathematically, connected with each other by means of geometric (conifold) transitions.*

This is the so called *Calabi–Yau web conjecture* described in many insightful papers starting from 1988. The word "mathematically" in the statement above is a prelude to the further problem of understanding how physics passes through the singularities of a geometric transition process. Actually, as far as I know, conifold transitions are almost the only g.t.s which have been understood from the physical point of view, after the work of A. Strominger [35]: the word "almost" refers to the so-called *hyperconifolds transitions*, which are divisorial g.t.s, discovered by R. Davies, having the property of being *mirror reverse transitions* of conifold ones (see [8] and [9]). Let us here underline that the hypothesis of simply connectedness is then necessary if one would only use conifold transitions, as a consequence of Corollary 3.4.

In this sense, conifold transitions turn out to be very interesting both mathematically and physically: probably because they give a concrete bridge between the *complex structures* and the *symplectic structures* on a Calabi–Yau 3-fold.

6. THE GROSS CALABI–YAU WEB: NODES AND ARROWS

A mathematically refined version of the Calabi–Yau web conjecture was presented by M. Gross in [18]: it is a sort of synthesis between the Reid's fantasy and the Calabi–Yau web.

The construction.

1. On the contrary of the K3 case for which an algebraic K3 surface can be smoothly deformed to a non–algebraic one, the deformation of a projective Calabi–Yau 3–fold, even singular, is still projective.
2. Since the hardest part of the Reid's fantasy seems to be in dealing with non–Kähler 3–folds, one could skip this part by insisting on staying within the projective category.
3. Think the nodes of the giant web predicted by the web conjecture as consisting of (suitable closures of) *def-equivalence classes* of Calabi–Yau 3–folds.
4. Two such nodes, say $\mathcal{M}_1$ and $\mathcal{M}_2$, are *connected* by an arrow $\mathcal{M}_1 \to \mathcal{M}_2$ if *there exist Calabi–Yau 3-folds* $Y \in \mathcal{M}_1$ *and* $\widetilde{Y} \in \mathcal{M}_2$ *which are each other connected by means of a geometric transition.* More precisely there exists:
   - a birational contraction to a normal 3–fold $\phi : Y \to \overline{Y}$
   - a deformation family $(\overline{\mathcal{Y}}, \overline{Y}) \to (\Delta, 0)$ such that $\overline{\mathcal{Y}}_t \cong \widetilde{Y}$ for some $t \in \Delta\ ,\ t \neq 0$.

Example 6.1 (See also [18]). Let

- $\mathcal{M}_Q$ be the family of quintic 3–folds in $\mathbb{P}^4$,
- $\mathcal{M}_D$ be the family of double solids (i.e. double covers of $\mathbb{P}^3$) branching along a octic surface of $\mathbb{P}^3$,
- $\mathcal{M}_T$ be (a closure of) the def-class of a smooth blow–up of a quintic 3–fold having a triple point.

Then these deformation families are nodes of the following connected graph

$$\begin{tikzcd}
 & \mathcal{M}_T \arrow[dl] \arrow[dr] & \\
\mathcal{M}_Q & & \mathcal{M}_D
\end{tikzcd} \tag{4}$$

where the two arrows are obtained as follows:

- let Z be a smooth element in $\mathcal{M}_T$ and $\phi : Z \to \overline{Y}$ be the contraction of the exceptional divisor of Z. Then $\overline{Y}$ is a quintic 3–fold in $\mathbb{P}^4$ with a triple point. Since $\overline{Y}$ can be smoothed to a quintic 3–fold we have $\mathcal{M}_T \longrightarrow \mathcal{M}_Q$
- if we project $\overline{Y}$ from its triple point then we get a rational morphism $\psi : \overline{Y} \dashrightarrow \mathbb{P}^3$ which can be lifted to the blow up Z of $\overline{Y}$, giving rise to a generically finite morphism $\widehat{\psi} : Z \to \mathbb{P}^3$. Consider its Stein factorization $\widehat{\psi} = f \circ \varphi$. Then we get the following commutative diagram

$$\begin{tikzcd}
Z \arrow[r, "\varphi"] \arrow[d, "\phi"] \arrow[dr, "\widehat{\psi}"] & \overline{X} \arrow[d, "f"] \\
\overline{Y} \arrow[r, dashed, "\psi"] & \mathbb{P}^3
\end{tikzcd} \tag{5}$$

  where f gives to $\overline{X}$ the structure of a double solid branched along a singular octic surface $S \subset \mathbb{P}^3$. Since $\overline{X}$ can immediately be smoothed by smoothing the branching locus $S \subset \mathbb{P}^3$ this gives the arrow $\mathcal{M}_T \longrightarrow \mathcal{M}_D$.

Conjecture 6.2 (of Connectedness)**.** *The graph of (simply connected) Calabi–Yau 3–folds is connected. Then their moduli can be described by starting from the* primitive *nodes given by def-classes of Calabi–Yau 3-folds which do not admits any* *birational contraction landing to a projective normal 3-fold (in general those having Picard number 1).*

A major evidence for this conjecture is given by Chiang-Greene-Gross-Kanter in [6] where the authors announced that, by computer procedure, *it is possible to settle in a connected graph all known examples of Calabi–Yau hypersurfaces in a 4-dimensional weighted projective space (7555 Calabi–Yau 3–folds).* Actually this web is even bigger since many weighted hypersurfaces has been connected passing through hypersurfaces of more general toric, Fano 4-dimensional varieties.
Moreover, lot of arrows in the previous big connected graph are not generated by conifold transitions but they are represented by very general geometric transitions. In fact many nodes of such a big connected graph are def-classes of non simply connected Calabi–Yau 3-folds, allowing us to drop the simply connectedness hypothesis in the statement of the Connectedness Conjecture 6.2.

7. Deformation of a morphism

Let $\phi : Y \to X$ be a morphism of complex spaces and let B be a connected complex space with a special point $o \in B$ such that $g : (\mathcal{Y}, Y) \to (B, o)$ and $f : (\mathcal{X}, X) \to (B, o)$ are deformation families of Y and X, respectively. Then a *deformation family of the morphism* ϕ is a morphism $\Phi : \mathcal{Y} \to \mathcal{X}$ such that the following diagram commutes

(6)
$$\begin{tikzcd}
Y \arrow[rrr, hook] \arrow[dr, "\phi"] \arrow[ddr] & & & \mathcal{Y} \arrow[dl, "\Phi"'] \arrow[ddl, "g"] \\
& X \arrow[r, hook] \arrow[d] & \mathcal{X} \arrow[d, "f"'] & \\
& o \arrow[r, phantom, "\in"] & B &
\end{tikzcd}$$

with $Y = \overline{g}^{-1}(o)$, $X = f^{-1}(o)$ and $\phi = \Phi_{|g^{-1}(o)}$.
Given two distinct points $b_1, b_2 \in B$ the morphism $\phi_2 := \Phi_{|g^{-1}(b_2)}$ is called a *deformation* of the morphism $\phi_1 := \Phi_{|g^{-1}(b_1)}$ and viceversa.
Let us now introduce a non-standard notation: the morphism deformation family $\mathcal{Y} \xrightarrow{\Phi} \mathcal{X}$ is called *smooth* if the deformation family $\mathcal{Y} \xrightarrow{g} B$ is smooth. In this case ϕ_2 will be called a *smooth* deformation of ϕ_1 and viceversa.

7.1. Deformation equivalence of geometric transitions.

Definition 7.1. Two g.t.s $T_1(Y_1, \overline{Y}_1, \widetilde{Y}_1)$ and $T_2(Y_2, \overline{Y}_2, \widetilde{Y}_2)$ are *deformation equivalent* (or *def-equivalent*) if

(1) $Y_1 \sim Y_2$,
(2) $\widetilde{Y}_1 \sim \widetilde{Y}_2$,
(3) the associated birational contractions $\phi_1 : Y_1 \to \overline{Y}_1$ and $\phi_2 : Y_2 \to \overline{Y}_2$ are deformations of each other.

We will write $T_1 \sim T_2$. This gives an equivalence relation on the set of geometric transitions.

Remark 7.2. Notice that *def-equivalent transitions* connect the same def-equivalence classes of Calabi–Yau 3-folds (i.e. the same nodes of the Gross Calabi–Yau web). As an immediate consequence:

- *an arrow of the Gross Calabi–Yau web is actually a def-equivalence class of geometric transitions.*

By stressing this concept we get that

- *the Gross Calabi–Yau web is a sort of quotient, up to def equivalence, of the string theoretic Calabi–Yau web restricted to the algebraic category.*

The previous Definition 7.1 can be simplified by putting hypothesis on the singular loci of def-equivalent g.t.s:

Proposition 7.3. *Let* $T_1(Y_1, \overline{Y}_1, \widetilde{Y}_1)$ *and* $T_2(Y_2, \overline{Y}_2, \widetilde{Y}_2)$ *be g.t.s such that both* $\mathrm{Sing}(\overline{Y}_1)$ *and* $\mathrm{Sing}(\overline{Y}_2)$ *comprise at most isolated terminal singularities. Then* $T_1 \sim T_2$ *if and only if their associated birational contractions* $\phi_i : Y_i \to \overline{Y}_i$ *are* smooth *deformations of each other.*

Proof. Assume that $T_1 \sim T_2$ then there exist two commutative diagrams

$$\begin{tikzcd}
Y_i \arrow[rrr, hook] \arrow[dr, "\phi"] \arrow[ddr] & & & \mathcal{Y} \arrow[dl, "\Phi"'] \arrow[ddl, "g"] \\
& \overline{Y}_i \arrow[r, hook] \arrow[d] & \overline{\mathcal{Y}} \arrow[d, "f"'] & \\
& b_i \arrow[r, phantom, "\in"] & B &
\end{tikzcd} \qquad (i = 1, 2)\ .$$

meaning that ϕ_1 and ϕ_2 are deformations of each other. Moreover $Y_1 \sim Y_2$ implies that the morphism deformation family Φ has to be smooth.
Viceversa if ϕ_1 and ϕ_2 are smooth deformations of each other then the deformation family $\mathcal{Y} \xrightarrow{g} B$ in the previous diagrams is necessarily smooth, giving $Y_1 \sim Y_2$. To prove that $\widetilde{Y}_1 \sim \widetilde{Y}_2$ observe that the generic fibre of the family $\overline{\mathcal{Y}} \xrightarrow{f} B$ admits at most isolated terminal singularities. Recall now the extension, due to Y. Namikawa, of the Bogomolov-Tian-Todorov Theorem 2.4 to the Kuranishi space of a terminal singularity, which turns out to be smooth ([23], Theorem 1). This means that, up to shrink it, the base space B may live on the closure of a unique def-equivalence class of Calabi–Yau threefolds. Since $\widetilde{Y}_i$ is a smoothing of $\overline{Y}_i$ this suffices to prove that both $\widetilde{Y}_1$ and $\widetilde{Y}_2$ have to belong to the same def-equivalence class. □

Let us say that hypothesis of the previous Proposition 7.3 are satisfied e.g. by *small* g.t.s (see the following Definition 8.6).

Remark 7.4. *The physics "passes through" def equivalence*, in the sense that it is *def-equivariant.*
In fact a conformal field theory on a Calabi–Yau 3-fold Y is the datum of a point χ on the *complexified Kähler cone*

$$\mathcal{K}_{\mathbb{C}}(Y) := \left\{ \chi \in H^2(Y, \mathbb{C}) \mid \Im(\chi) \in \mathcal{K}(Y) \right\} \big/ H^2(X, \mathbb{Z})$$

where $\mathcal{K}(Y)$ is the Kähler cone of Y and the action of integral cohomology is additive on the real part $\Re(\chi)$. In fact the imaginary part $\omega := \Im(\chi)$ gives the Kähler , Ricci flat metric of Y, which is the *curvature properties of the supersymmetric string theoretic vacuum*, while the real part $b := \Re(\chi)$ gives the so called *b-field* describing *the strings' charge properties* of the theory.

If $Y_1 \sim Y_2$ are two def equivalent Calabi–Yau 3-folds, then there exists an orientation preserving diffeomorphism $f : Y_1 \cong Y_2$ inducing a contravariant isomorphism $f^* : \mathcal{K}_{\mathbb{C}}(Y_2) \cong \mathcal{K}_{\mathbb{C}}(Y_1)$. Then the physical theories (Y_2, χ) and $(Y_1, f^*(\chi))$ are isomorphic.

8. Simple geometric transitions

Since both in mathematics and in physics the conifold transitions are the most understood geometric transitions between Calabi–Yau 3–folds, it makes sense to ask *when a geometric transition is def-equivalent to a conifold one.* Let us then set the following

Definition 8.1 (Simple g.t.s and arrows)**.** A g.t. is called *simple* if it is def-equivalent to a conifold transition. Therefore an arrow is called simple if it is the def-equivalence class of a conifold transition.

Remark 8.2 (The importance of being simple). If $T(Y, \overline{Y}, \widetilde{Y})$ is a simple g.t. then, for what observed above:

- it is physically well understood by the Remark 7.4,
- Y and $\widetilde{Y}$ are *almost diffeomorphic* i.e. there exist finite open coverings $\{U_i\}_{i=1}^N$ and $\{\widetilde{U}_i\}_{i=1}^N$ of Y and $\widetilde{Y}$, respectively, and almost everywhere defined diffeomorphisms from U_i to $\widetilde{U}_i$
$$\alpha_i : U_i \setminus \operatorname{Exc}(\phi) \xrightarrow{\cong} \widetilde{U}_i \setminus \mathcal{V} \ , \ 1 \leq i \leq N \ ,$$
where $\mathcal{V} \subset \widetilde{Y}$ is the vanishing locus. In particular Y and $\widetilde{Y}$ have the same fundamental group.

In fact if $T'(X, \overline{X}, \widetilde{X})$ is a conifold t. with $T \sim T'$, then there are orientation preserving diffeomorphisms $Y \cong X$ and $\widetilde{Y} \cong \widetilde{X}$. Moreover Theorem 3.3 ensures that X and $\widetilde{X}$ are almost diffeomorphic, then implying the same property for Y and $\widetilde{Y}$. In particular the cardinality N of the open coverings is that of the singular locus of $\overline{X}$, i.e. $N = |\operatorname{Sing}(\overline{X})|$. To end up, corollary 3.4 applies.

8.1. Type II geometric transitions are not simple.

Definition 8.3. A *type II geometric transition* $T(Y, \overline{Y}, \widetilde{Y})$ is a g.t. such that

- the associated birational morphism $\phi : Y \to \overline{Y}$ is *primitive*, i.e. it cannot be factored into birational morphisms of normal varieties,
- ϕ contracts a divisor down to a point; in this case the exceptional divisor E is irreducible and in particular it is a (generalized) *del Pezzo surface* (see [29]).

Example 8.4. The g.t. $T(Z, \overline{Y}, \widetilde{Y})$ representing the arrow $\mathcal{M}_T \longrightarrow \mathcal{M}_Q$ in the Example 6.1, is a type II g.t.

By exhibiting a suitable weighted blow down, one can easily produce a type II g.t. $T(Y, \overline{Y}, \widetilde{Y})$ such that Y and $\widetilde{Y}$ do not admit the same fundamental group. Then T *cannot be a simple g.t.* due to the previous Remark 8.2 and Corollary 3.4. Actually a much stronger result can be established:

Theorem 8.5. *A type II g.t. is never simple.*

Proof. Let us assume the existence of a def-equivalence between a type II g.t. $T_1(Y_1, \overline{Y}_1, \widetilde{Y}_1)$ and a conifold $T_2(Y_2, \overline{Y}_2, \widetilde{Y}_2)$, giving a deformation family of morphisms

$$\begin{tikzcd}
\mathcal{Y} \arrow[rr, "\Phi"] \arrow[dr, "f"'] & & \overline{\mathcal{Y}} \arrow[dl, "g"] \\
& B &
\end{tikzcd}$$

over a connected base B and such that, up to shrink the base, $\mathcal{Y} \stackrel{f}{\to} B$ is a smooth family realizing the def-equivalence $Y_1 \sim Y_2$. In particular there exist two distinct points $b_1, b_2 \in B$ such that $\Phi_{b_i} := \Phi_{|f^{-1}(b_i)} = \phi_i : Y_i \to \overline{Y}_i$. Since T_2 is conifold then $\mathrm{Exc}(\phi_2)$ is composed by a finite number of disjoint smooth rational curves whose normal bundle is given by $\mathcal{O}_{\mathbb{P}^1}(-1) \oplus \mathcal{O}_{\mathbb{P}^1}(-1)$ (so called $(-1,-1)$-*curves*). Then any $(-1,-1)$-curve is a *stable submanifold* of Y_2 in the sense of Kodaira [19]. For this reason, up to shrink the connected base B, we may now assume B to be a suitable neighborhood B° of b_2 with the addition of a closure point given by b_1, such that $\Phi_b : Y_b \to \overline{Y}_b$ turns out to be a c.t. for any $b \in B$ with $b \neq b_1$. The contradiction is then reached by observing that $\overline{Y}_1$ is $\mathbb{Q}$-factorial while $\overline{Y}_b$ can never be $\mathbb{Q}$-factorial for any $b \neq b_1$: this fact is against a result of J. Kollár and S. Mori ([21] Thm. (12.1.10)) guaranteeing that $\mathbb{Q}$-factoriality of the fibers has to be an *open condition* for the induced deformation family $g : \overline{\mathcal{Y}} \to B$ i.e. that there should exist an open neighborhood of $b_1 \in B$ over which all fibers should be $\mathbb{Q}$-factorial.

8.2. An example of a non-simple small geometric transition.

Definition 8.6 (Small g.t.). A g.t. $T(Y, \overline{Y}, \widetilde{Y})$ is called *small* if the associated birational morphism $\phi : Y \to \overline{Y}$ is a *small birational contraction*, i.e. its exceptional locus $\mathrm{Exc}(\phi)$ has codimension greater than 1 in Y.

Possible exceptional and singular loci occurring in a small g.t. are completely classified (see [32], Thm. 6 and references therein):

- $\mathrm{Sing}(\overline{Y})$ turns out to be composed by a finite number of isolated *compound Du Val (cDV)* singular points, which in particular are terminal singularities,
- $\mathrm{Exc}(\phi)$ is then composed by a finite number of trees of transversally intersecting rational curves, dually represented by ADE Dynkin graphs.

Due to the particular geometry of the exceptional locus $\mathrm{Exc}(\phi)$ it is quite natural to ask for the *simplicity of any small g.t.s.* Unfortunately this is not the case, as the following example shows.

Example 8.7. The following example is essentially due to Y. Namikawa ([24], Example 1.11).

Let S be the rational elliptic surface with sections obtained as the Weierstrass fibration associated with the bundles homomorphism

$$(0,B) \quad : \quad \mathcal{E} = \mathcal{O}_{\mathbb{P}^1}(3) \oplus \mathcal{O}_{\mathbb{P}^1}(2) \oplus \mathcal{O}_{\mathbb{P}^1} \longrightarrow \mathcal{O}_{\mathbb{P}^1}(6) \tag{7}$$
$$(x,y,z) \longmapsto -x^2 z + y^3 + B(\lambda)\ z^3$$

for a generic $B \in H^0(\mathbb{P}^1, \mathcal{O}_{\mathbb{P}^1}(6))$ i.e. S is the zero locus of $(0,B)$ in the projectivized bundle $\mathbb{P}(\mathcal{E})$. Then:

(1) *the natural fibration $S \to \mathbb{P}^1$ has generic smooth fibre and 6 distinct cuspidal fibres,*

(2) *the fiber product* $X := S \times_{\mathbb{P}^1} S$ *is a threefold admitting 6 singularities of type* $II \times II$*, in the standard Kodaira notation* [20],

(3) X *admits a* small *resolution* $\widehat{X} \xrightarrow{\phi} X$ *whose exceptional locus is composed by 6 disjoint couples of rational curves intersecting in one point i.e. 6 disjoint* A_2 *exceptional trees,*

(4) *by results of C. Schoen,* X *is a special fibre of the family of fiber products* $S_1 \times_{\mathbb{P}^1} S_2$ *of rational elliptic surfaces with sections: in particular for* S_1 *and* S_2 *sufficiently general* $\widetilde{X} = S_1 \times_{\mathbb{P}^1} S_2$ *is a Calabi–Yau threefold giving a smoothing of* X ([34] §2).

Since ϕ is a small, crepant resolution, $\widehat{X}$ turns out to be a Calabi–Yau threefold and $T(\widehat{X}, X, \widetilde{X})$ is a small *non–conifold* g.t.. Let p be one of the six singular points of X, locally defined as a germ of singularity by the polynomial

$$F := x^2 - z^2 - y^3 + w^3 \in \mathbb{C}[x, y, z, w] \ .$$

Consider the *localization near to* p

$$\begin{tikzcd} \widehat{U}_p := \phi^{-1}(U_p) \arrow[r, hook] \arrow[d, "\varphi"] & \widehat{X} \arrow[d, "\phi"] \\ U_p := \operatorname{Spec} \mathcal{O}_{F,p} \arrow[r, hook] & X \end{tikzcd} \tag{8}$$

which induces, since p is a rational singularity, the following commutative diagram of maps between Kuranishi spaces

$$\begin{tikzcd} \operatorname{Def}(\widehat{X}) \arrow[r, "\widehat{l}_p"] \arrow[d, hook, "D"] & \operatorname{Def}(\widehat{U}_p) \arrow[d, hook, "D_{loc}"] \\ \operatorname{Def}(X) \arrow[r, "l_p"] & \operatorname{Def}(U_p) \cong T^1_{U_p} \end{tikzcd} \tag{9}$$

where the horizontal maps are the natural localization maps while the vertical maps are *injective maps* induced by the resolution ϕ (see [38] Propositions 1.8 and 1.12, [21] Proposition (11.4)). Then, by explicit calculations (see [**?**], Thm. 3.6), it turns out that

$$\left.\begin{array}{c} \dim \operatorname{Def}(\widehat{U}_p) = 1 \\ \operatorname{Im}(l_p) \cap \operatorname{Im}(D_{loc}) = 0 \end{array}\right\} \Longrightarrow \operatorname{Im}(\widehat{l}_p) = 0 \tag{10}$$

meaning that

(5) *no global deformation of* $\widehat{X}$ *may induce a local non–trivial deformation of* $\widehat{U}_p$*; in particular* $\phi^{-1}(p)$ *turns out to be a* rigid A_2 *exceptional tree and* $T(\widehat{X}, X, \widetilde{X})$ *cannot be def-equivalent to a conifold transition.*

8.3. **A cohomological condition for simplicity of small transitions.** Anyway the previous example allows us to understand some further conditions that a small g.t. should satisfy to be a simple g.t.:

Theorem 8.8. *Let* $\Theta_\bullet$ *be the dual sheaf of the Kähler differentials sheaf* $\Omega_\bullet$*. Then if* $T(Y, \overline{Y}, \widetilde{Y})$ *is a* general *small geometric transition such that*

$$h^1(\overline{Y}, \Theta_{\overline{Y}}) < h^1(Y, \Theta_Y) \tag{11}$$

then T *is a simple g.t..*

Sketch of proof. The proof is an application of R. Friedman techniques presented in [12]. In fact by the Leray spectral sequence applied to the birational small contraction $\phi : Y \to \overline{Y}$ and the local to global spectral sequence relating the global $\mathbb{T}^{\bullet}_{\overline{Y}} := \mathrm{Ext}^{\bullet}(\Omega_{\overline{Y}}, \mathcal{O}_{\overline{Y}})$ with the sheaves $\Theta^{\bullet}_{\overline{Y}} := \mathcal{E}xt^{\bullet}(\Omega_{\overline{Y}}, \mathcal{O}_{\overline{Y}})$, one gets the following commutative diagram

$$\begin{tikzcd}
0 \to H^1(Y, R^0\phi_*\Theta_Y) \arrow[r] \arrow[d, equal] & \mathbb{T}^1_Y \arrow[r, "\lambda"] \arrow[d, "\delta"] & H^0\left(\overline{Y}, R^1\phi_*\Theta_Y\right) \arrow[r] \arrow[d, "\delta_{loc}"] & \cdots \\
0 \to H^1(\overline{Y}, \Theta_{\overline{Y}}) \arrow[r] & \mathbb{T}^1_{\overline{Y}} \arrow[r, "\overline{\lambda}"] & T^1_{\overline{Y}} := H^0(\overline{Y}, \Theta^1_{\overline{Y}}) \arrow[r] & \cdots
\end{tikzcd} \tag{12}$$

where

- the vertical equality come from an application of Hartogs Theorem giving $R^0\phi_*\Theta_Y \cong \Theta_{\overline{Y}}$ ([12] Lemma (3.1)),
- the vertical morphism δ is the differential of an injective map between Kuranishi spaces $\mathrm{Def}(Y) \hookrightarrow \mathrm{Def}(\overline{Y})$, constructed by J. Kollar and S. Mori ([21] Proposition (11.4)), which turns out to be still injective since δ_{loc} is injective,
- δ_{loc} is the localization of δ near to $\mathrm{Sing}(\overline{Y})$, which is injective by a result of Friedman ([12], Prop. (2.1)).

Since $\mathbb{T}^1_Y \cong H^1(Y, \Theta_Y)$, a *first meaning of the generality hypothesis* in the statement is the request that the cohomological condition (11) would guarantee the existence of a global first order deformation ξ of Y inducing a non-trivial local deformation $\xi_{loc} := \delta_{loc} \circ \lambda(\xi)$ giving non trivial first order deformation of *any* singularity $p \in \mathrm{Sing}(\overline{Y})$: this means that $\mathrm{Im}\,\lambda$ has to be represented by a point in general position inside a suitable grassmannian of subspaces of $H^0\left(\overline{Y}, R^1\phi_*\Theta_Y\right)$. Since p is an isolated cDV point, its Kuranishi space is smooth and well described. In particular the locus of local first order deformations sending p to an ordinary double point, describes an hypersurface $\mathcal{N}_p$ in the tangent space $H^1(\overline{U}_p, \Theta_{\overline{U}_p})$ to the Kuranishi space of a neighborhood $\overline{U}_p \subset \overline{Y}$ of p.
A second level of generality for T resides in requiring that a *sufficiently general* $\xi \in \mathbb{T}^1_Y$ generates, for any $p \in \mathrm{Sing}(\overline{Y})$, a pencil inside $\mathcal{N}_p$ whose general fiber has singular locus comprising ordinary double points only. The following Example 8.4 shows that such a general g.t. can actually occur.
For full details the interested reader is referred to the forthcoming [33]. □

Remark 8.9. Back to the Namikawa's example 8.7, let us observe that for the g.t. $T(\widehat{X}, X, \widetilde{X})$, with $X = S \times_{\mathbb{P}^1} S$ be the fibred self-product of a cuspidal elliptic surface, one gets $h^1(\overline{Y}, \Theta_{\overline{Y}}) = h^1(Y, \Theta_Y) = 3$.

8.4. **An example of a simple small geometric transition.** Let us consider the singular quintic threefold $\overline{Q} \subset \mathbb{P}^4$ given by

$$u(u-2x)(u-3y)(x^2-y^2) - (z^5 - w^5) = 0\ . \tag{13}$$

The singular locus $\mathrm{Sing}(\overline{Q})$ is composed by 10 isolated hypersurface singularities, each of them analytically equivalent to the one described by the local equation

$$x^2 - y^2 = z^5 - w^5 \tag{14}$$

which is a cA_4 singular point whose Milnor and Tyurina numbers are equal to 16. A resolution of this singular point is obtained by a *successive blow up* of the planes

$$\pi_i : x - y = z - \epsilon^i w = 0 \quad , \quad 0 \leq i \leq 3 \ , \ \epsilon^5 = 1 \ .$$

More precisely: blow up $\mathbb{C}^4$ along π_0, then blow up the obtained $\mathbb{C}^5$ along the strict transform of π_1 and so on. At the end look at the strict transform of the singularity, which carries an exceptional locus composed by a tree of 4 lines dually represented by the Dynkin graph A_4.

We are now in a position to construct a non-conifold geometric transition as follows:

- *the resolution:* the quintic threefold $\overline{Q}$ admits a global resolution $\widehat{Q}$ which can be obtained by the successive blow up of 16 planes

  $$\pi_i^j : l_j = z - \epsilon^i w = 0 \quad 0 \leq i \leq 3 \ , \ 1 \leq j \leq 4$$

  where $\{l_1, \ldots, l_4\} \subset \{u, u - 2x, u - 3y, x - y, x + y\}$.
- *the smoothing:* it is obviously given by a smooth quintic threefold $Q \subset \mathbb{P}^4$.

This gives the g.t. $T(\widehat{Q}, \overline{Q}, Q)$. To deform T to a conifold transition consider the following deformation $\overline{Q}_{(a,b,c)}$ of $\overline{Q}$

$$u(u-2x)(u-3y)(x^2-y^2)-(z-w)(z-\epsilon w)(z-\epsilon^2 w+a)(z-\epsilon^3 w+b)(z-\epsilon^4 w+c)=0$$

which, for a general $\alpha := (a, b, c) \in \mathbb{C}^3$, splits up each singular point of $\overline{Q}$ into 10 nodes, hence giving 100 nodes. Since the deformation $\overline{Q}_\alpha$ respects the factorization in the equation of $\overline{Q}$, it lifts to a deformation $\widehat{Q}_\alpha$ of the resolution $\widehat{Q}$ splitting up every exceptional A_4 tree into 10 disjoint lines. This gives a deformation family of morphisms

$$\begin{tikzcd}
\widehat{\mathcal{Q}} \arrow[rr, "\Phi"] \arrow[dr, "f"'] & & \overline{\mathcal{Q}} \arrow[dl, "g"] \\
& \mathbb{C}^3 &
\end{tikzcd}$$

hence a def-equivalence $T \sim T_\alpha(\widehat{Q}_\alpha, \overline{Q}_\alpha, Q)$. Let us further observe that the deformations $\overline{Q}_\alpha$ are not all distinct up to isomorphisms: if we consider the Kuranishi space T^1 of any singularity of $\overline{Q}$, there is a well defined map $\mathbb{C}^3 \to T^1$ whose image is 1-dimensional. This is enough to show that

$$\dim \left(\operatorname{Im} \left(\lambda : \mathbb{T}^1_{\widehat{Q}} \to H^0(\overline{Q}, R^1\phi_*\Theta_{\widehat{Q}}) \right) \right) = 1$$

giving $h^1(\Theta_{\overline{Q}}) = 17 < 18 = h^1(\Theta_{\widehat{Q}})$. The further main invariants of the g.t. T and the conifold t. T_α are listed in the following table:

Variety	$h^1(\Theta_\bullet)$	b_2	ρ	b_3	b_4	χ
$\widehat{Q}, \widehat{Q}_\alpha$	18	17	17	38	17	-2
$\overline{Q}$	17	1	1	62	17	-42
$\overline{Q}_\alpha$	18	1	1	122	17	-102
Q	101	1	1	204	1	-200

They can be computed from the well known invariants of the smooth quintic threefold Q by means of relations given in [32], Thm. 7.

References

[1] Batyrev V. "Dual polyhedra and mirror symmetry for Calabi–Yau hypersurfaces in toric varieties" *J. Alg. Geom.* **3** (1994), 493–535; `math.AG/9310003`.

[2] Bauer I. C., Catanese F. and Pignatelli R. "Complex surfaces of general type: some recent progress" *Global aspects of complex geometry*, 1–58, Springer, Berlin, 2006.

[3] Bogomolov F. "Hamiltonian Kähler manifolds" *Dokl.Akad.Nauk.SSSR* **243/5** (1978), 1101–1104.

[4] Catanese F. "Moduli spaces of surfaces and real structures" *Ann. of Math.* **158** (2003), 577–592.

[5] Catanese F. and Wajnryb B. "Diffeomorphism of simply connected algebraic surfaces" *J. Differential Geom.* **76** (2007), 177213.

[6] Chiang T.M., Greene B., Gross M. and Kanter Y. "Black Hole condensation and the web of Calabi–Yau manifolds" *Nucl.Phys.Proc.Suppl.* **46** (1996), 82–95; `hep-th/9511204`.

[7] Clemens C.H. "Double Solids" *Adv. in Math.* **47** (1983), 107–230.

[8] Davies R. "Quotients of the conifold in compact Calabi-Yau threefolds and new topological transitions" *Adv. Theor. Math. Phys.* **14** (2010), 965–989; `arXiv:0911.0708`.

[9] Davies R. "Hyperconifold transitions, mirror symmetry, and string theory" *Nuclear Phys. B* **850** (2011), 214–231; `arXiv:1102.1428`.

[10] Douady A. "Le problème des modules locaux pour les espaces C–analytiques compacts" *Ann. scient. Éc. Norm. Sup.* 4e série, **7** 569–602 (1974).

[11] Enriques F. *Le superficie algebriche* Zanichelli, Bologna (1946).

[12] Friedman R. "Simultaneous resolution of threefold double points" *Math. Ann.* **247** (1986), 671–689.

[13] Friedman R. *Donaldson and Seiberg-Witten invariants of algebraic surfaces*, Algebraic geometrySanta Cruz 1995, 85–100, *Proc. Sympos. Pure Math.*, **62**, Amer. Math. Soc., Providence, RI, 1997.

[14] Friedman R. and Morgan J. W. "Algebraic surfaces and 4-manifolds: some conjectures and speculations" *Bull. Amer. Math. Soc.* (N.S.) **18** (1988), 1–19.

[15] Grassi A. and Rossi M. "Large N–dualities and transitions in geometry" in *Geometry and Physics of Branes, Como 2001* Series of H.E.P., Cosmology and Gravitation, IoP Bristol (2003), 210–278; `math.AG/0209044`.

[16] Grauert H. "DerSatz von Kuranishi für Kompakte Komplexe Räume" *Invent.Math.* **25**, 107–142 (1974).

[17] Greene B., Morrison D.R. and Strominger A. "Black hole condensation and the unification of string vacua" *Nucl. Phys.* **B451** (1995), 109–120; `hep-th/9504145`.

[18] Gross M. "Primitive Calabi–Yau threefolds" *J. Diff. Geom.* **45** (1997), 288–318; `math.AG/9512002`.

[19] Kodaira K. "On stability of compact submanifolds of complex manifolds" *Amer. J. Math.* **85** (1963), 79–94.

[20] Kodaira K. "On the structure of compact complex analytic surfaces" *Am. J. Math.* **86** (1964), 751–798.

[21] Kollár J. and Mori S. "Classification of three–dimensional flips" *J. Amer. Math. Soc.* **5** (1992), 533–703.

[22] Manetti M. "On the moduli space of diffeomorphic algebraic surfaces" *Invent. Math.* **143** (2001), 29–76.

[23] Namikawa Y. "On deformations of Calabi-Yau 3-folds with terminal singularities" *Topology* **33**(3) (1994), 429–446.

[24] Namikawa Y. "Stratified local moduli of Calabi–Yau 3–folds" *Topology* **41** (2002), 1219–1237.

[25] Palamodov V. P. "The existence of versal deformations of complex spaces" *Dokl. Akad. Nauk SSSR* **206** (1972), 538–541.

[26] Palamodov V. P. "Deformations of complex spaces" *Russian Math. Surveys* **31**(3) (1976), 129–197; from russian *Uspekhi Mat. Nauk* **31**(3) (1976), 129–194.

[27] Ran Z. "Deformations of maps" in *Algebraic curves and projective geometry*, Ballico E. and Ciliberto C., Eds. LNM **1389**, Springer–Verlag (1989).

[28] Ran Z. "Deformations of manifolds with torsion or negative canonical bundle" *J.Alg.Geom.* **1** (1992), 279–291.

[29] Reid M. "Canonical 3–folds" in *Journées de géométrie algébrique d'Angers*, Sijthoff & Nordhoff (1980), 671–689.

[30] Reid M. "The moduli space of 3–folds with $K = 0$ may neverthless be irreducible" *Math. Ann.* **287** (1987), 329–334.

[31] Rossi M. "Geometric transitions" *J. Geom. Phys.* **56**(9) (2006), 1940–1983.

[32] Rossi M. "Homological type of geometric transitions" *Geom. Dedicata* **151** (2011), 323–359.

[33] Rossi M. "Def-equivalence of geometric transitions" in preparetion.

[34] Schoen C. "On fiber products of rational elliptic surfaces with section" *Math. Z.* **197**(2) (1988), 177–199.

[35] Strominger A. "Massless black holes and conifolds in string theory" *Nucl. Phys.* **B451** (1995), 97–109; `hep-th/9504145`.

[36] Tian G. "Smoothness of the universal deformation space of compact Calabi–Yau manifolds and its Weil–Petersson metric" in *Mathematical aspects of string theory* (S.-T. Yau, ed.) World Scientific, Singapore (1987), 629–646.

[37] Todorov A. "The Weil-Petersson geometry of the moduli space of SU($n \geq 3$) (Calabi–Yau) manifolds" *Comm.Math.Phys.* **126** (1989), 325–346.

[38] Wahl, J.M. "Equisingular deformations of normal surface singularities, I" *Ann. of Math.* **104** (1976), 325–356.

[39] Wilson P.M.H. "The Kähler cone on Calabi–Yau threefolds" *Invent.Math.* **107** (1992), 561–583.

[40] Wall C.T.C. "Classification problems in topology V: on certain 6–manifolds" *Invent.Math.* **1** (1966), 355-374.

Dipartimento di Matematica, Università di Torino, via Carlo Alberto 10, 10123 Torino

E-mail address: `michele.rossi@unito.it`